COMP60121: Automated Reasoning, Part II
Advanced Topics

Renate Schmidt

School of Computer Science
University of Manchester

schmidt@cs.man.ac.uk

http://www.cs.man.ac.uk/~schmidt/COMP6012/

Overview

- Handouts: These notes
  Weekly assessed coursework exercises
  Weekly unassessed exercises
  Using SPASS (distributed next week)
  Timetable (note: approximate!)
  Exam definition sheet (let me know of any mistakes)
- Deadlines for assessed coursework: 9:00 on next teaching day
  see website
- Demonstrators: Konstantin Korovin
  Nestan Tsiskaridze
  Adam Trybus
- Closed book exam

Content

- Emphasis on:
  - foundation of advanced automated theorem proving
  - both theory and practice (more theoretical, but motivated by considerations of practical aspects and efficiency)
- Treatment is formal and rigorous; small selection of important topics; many examples and exercises
- **All you need to know for the exam is in the lecture notes, and it is advisable that you can solve the exercises and can do the assignments**
- Good strategy: go through the material discussed in lectures after each lecture, do as many of the unassessed exercises, do the assignments and read ahead. And, ask questions!

Textbooks

- None cover all the material, but recommended are:
  - Schöning, U. (1989), *Logic for Computer Scientists.* Birkhäuser.
  - Fitting, M. (1990), *First-Order Logic and Automated Theorem Proving.* Springer.
- Also useful:
  - Socher-Ambrosius, R., Johann, P. (1997), *Deduction Systems.* Springer.
  - Goubault-Larrecq, J. and Mackie, I. (1997), *Proof Theory and Automated Deduction.* Kluwer.
  - Leitsch, A. (1997), *The Resolution Calculus.* Springer.
- All available in the Resources Centre Library and/or the main library. There is no need to buy a book.

COMP60121: Automated Reasoning II

Lecture 1

Previously . . . (in Part I)

- Syntax and semantics of propositional logic and FOL
  - variables, constants, terms, quantifiers, . . .
- Syntax and semantics of clause logic
  - atoms, literals, clauses, Skolem terms
- Transformation of formulae to clausal form
- Resolution calculus
  - resolution rule
  - factoring rule
  - unification – for first-order clauses

Aims

- To discuss and study optimised transformations into clausal form
- To introduce well-founded orderings
- To introduce ordered resolution with (well-behaved!) selection
- To study and show soundness and refutational completeness of resolution for ground case
- To introduce redundancy elimination
- To discuss applications

Well-Founded Orderings

- To show the refutational completeness of resolution, we make use of the concept of well-founded orderings.
- They are essential for making the search more deterministic and efficient.
- Are used in refinements of resolution.
- Reference:
  Baader, F. and Nipkow, T. (1998), *Term rewriting and all that*. Cambridge Univ. Press, Chapter 2.

Basic Properties of Relations

Let R be a binary relation over a set X ($R \subseteq X \times X$).

- R is transitive iff
  $\forall x, y, z \in X$, if $R(x, y)$ and $R(y, z)$ then $R(x, z)$.
- R is irreflexive iff $\forall x \in X$, $\neg R(x, x)$
- R is total, or linear, iff
  $\forall x, y \in X$, if $x \neq y$ then $R(x, y)$ or $R(y, x)$
- R^* denotes the reflexive-transitive closure of R. I.e.

$$R^* = \bigcup_{n=0}^{\infty} R^n = R^0 \cup R \cup R^2 \cup R^3 \cup \ldots$$

  where $R^0(x, x)$ for any $x \in X$ and
  $R^{n+1}(x, y)$ iff $\exists z.\ R(x, z) \wedge R^n(z, y)$ for $n \geq 0$.

Orderings

- A (strict) ordering on a set X is a transitive and irreflexive binary relation on X, here denoted by $\succ$.
- The pair $(X, \succ)$ is then called a (strictly) ordered set.
- An element x of X is minimal wrt. $\succ$, if there is no y in X such that $x \succ y$.
- A minimal element x in X is called the smallest (or strictly minimal) element, if for all $y \in X$ different from x, $y \succ x$.
- Maximal and largest (or strictly maximal) elements are defined analogously.
- **Notation**: $\prec$ for the inverse relation $\succ^{-1}$
  $\succeq$ for the reflexive closure ($\succ \cup =$) of $\succ$, i.e.
  $x \succeq y$ iff either $x \succ y$ or $x = y$

Well-Foundedness

- A (strict) ordering $\succ$ over X is called well-founded (or Noetherian or terminating), if there is no infinite decreasing chain $x_0 \succ x_1 \succ x_2 \succ \ldots$ of elements $x_i \in X$.

Lemma 1

$(X, \succ)$ is well-founded iff every non-empty subset Y of X has a minimal element.

- **Examples:**
  - Natural numbers: $(\mathbb{N}, >)$
- **Counterexamples:**
  - $(\mathbb{Z}, >)$
  - $(\mathbb{N}, <)$

Noetherian Induction (optional)

Property 2 (Noetherian Induction)

Let $(X, \succ)$ be a well-founded ordering, let Q be a property of elements of X.

If for all $x \in X$ the following implication is satisfied

if $Q(y)$ holds, for all $y \in X$ such that $x \succ y$, [a]
then $Q(x)$ holds. [b]

then

the property $Q(x)$ holds for all $x \in X$.

[a] induction hypothesis
[b] induction step

Noetherian Induction (optional) (cont'd)

Proof: By contradiction.
Thus, suppose for all $x \in X$ the implication above is satisfied, but $Q(x)$ does not hold for all $x \in X$.
Let $A = \{x \in X \mid Q(x) \text{ is false}\}$. Suppose $A \neq \emptyset$.
Since $(X, \succ)$ is well-founded, A has a minimal element x_1. Hence for all $y \in X$ with $x_1 \succ y$ the property $Q(y)$ holds.
On the other hand, the implication which is presupposed for this theorem holds in particular also for x_1, hence $Q(x_1)$ must be true so that x_1 cannot belong to A. *Contradiction*.

Multi-Sets

- Multi-sets are "sets which allow repetition".
  E.g.: $\{a, a, b\}$, $\{a, b, a\}$, $\{a, b\}$
- Formally, let X be a set.
  A multi-set S over X is a mapping $S : X \longrightarrow \mathbb{N}$.
- Intuitively, $S(x)$ specifies the number of occurrences of the element x (of the base set X) within S.
- We say that x is an element of S, if $S(x) > 0$.
- We use set notation ($\in$, $\subset$, $\subseteq$, $\cup$, $\cap$, etc.) with analogous meaning also for multi-sets, e.g.,

$$\begin{aligned}(S_1 \cup S_2)(x) &= S_1(x) + S_2(x)\\ (S_1 \cap S_2)(x) &= \min\{S_1(x), S_2(x)\}\end{aligned}$$

Multi-Sets (cont'd)

- **Example:** $S = \{a, a, a, b, b\}$ is a multi-set over $\{a, b, c\}$, where $S(a) = 3$, $S(b) = 2$, $S(c) = 0$.
- A multi-set S over X is called finite, if

$$|\{x \in X \mid S(x) > 0\}| < \infty.$$

- **From now on we consider finite multi-sets only.**

Exercise

Suppose $S_1 = \{c, a, b\}$ and $S_2 = \{a, b, b, a\}$ are multi-sets over $\{a, b, c, d\}$.
Determine $S_1 \cup S_2$ and $S_1 \cap S_2$.

Exercise

Suppose $S_1 = \{c, a, b\}$ and $S_2 = \{a, b, b, a\}$ are multi-sets over $\{a, b, c, d\}$.
Determine $S_1 \cup S_2$ and $S_1 \cap S_2$.

Answer:

$$S_1 \cup S_2 = \{a, a, a, b, b, b, c\}$$
$$S_1 \cap S_2 = \{a, b\}$$

Multi-Set Orderings

- Let $(X, \succ)$ be an ordering. The multi-set extension $\succ_{\text{mul}}$ of $\succ$ to (finite) multi-sets over X is defined by

$$S_1 \succ_{\text{mul}} S_2 \;\text{ iff }\; S_1 \neq S_2 \text{ and } \forall x \in X, \text{ if } S_2(x) > S_1(x) \text{ then } \exists y \in X : \; y \succ x \text{ and } S_1(y) > S_2(y)$$

- **Example:** Over $(\mathbb{N}, >)$:

$$\{5, 5, 4, 3, 2\} >_{\text{mul}} \{5, 4, 4, 3, 3, 2\} >_{\text{mul}} \{5, 4, 3\}$$

Exercise: How is the set $\{5, 5\}$ related to each of these?

Method for Determining $\succ_{\text{mul}}$

1. Remove common occurrences of elements from S_1 and S_2. Assume this gives S_1' and S_2'.
2. Then check that for every element x in S_2' there is an element $y \in S_1'$ that is larger than x. Then $S_1 \succ_{\text{mul}} S_2$.

- This method facilitates this equivalent definition:

$$S_1 \succ_{\text{mul}} S_2 \;\text{ iff }\; S_1 \neq S_2 \text{ and } \forall x \in S_2 \backslash S_1.\ \exists y \in S_1 \backslash S_2.\ y \succ x$$

Reconsider Example

- $S_1 = \{\cancel{5}, 5, \cancel{4}, \cancel{3}, \cancel{2}\}$ $\quad S_2 = \{\cancel{5}, \cancel{4}, 4, \cancel{3}, 3, \cancel{2}\}$
  $S_1' = \{5\}$ $\quad S_2' = \{4, 3\}$
  $5 > 4$ and $5 > 3$
  Therefore $S_1 >_{\text{mul}} S_2$.
- $S_2 = \{\cancel{5}, \cancel{4}, 4, \cancel{3}, 3, 2\}$ $\quad S_3 = \{\cancel{5}, \cancel{4}, \cancel{3}\}$
  $S_2' = \{4, 3, 2\}$ $\quad S_3' = \emptyset$
  Therefore $S_2 >_{\text{mul}} S_3$.
- Exercise: How does $S_4 = \{5, 3, 2\}$ compare with S_3?

Reconsider Example

- $S_1 = \{\not{B}, 5, \not{A}, \not{B}, \not{2}\}$ $\quad S_2 = \{\not{B}, \not{A}, 4, \not{B}, 3, \not{2}\}$
  $S_1' = \{5\}$ $\quad S_2' = \{4, 3\}$
  $5 > 4$ and $5 > 3$
  Therefore $S_1 >_{\text{mul}} S_2$.
- $S_2 = \{\not{B}, \not{A}, 4, \not{B}, 3, 2\}$ $\quad S_3 = \{\not{B}, \not{A}, \not{B}\}$
  $S_2' = \{4, 3, 2\}$ $\quad S_3' = \emptyset$
  Therefore $S_2 >_{\text{mul}} S_3$.
- Exercise: How does $S_4 = \{5, 3, 2\}$ compare with S_3?

  Answer: $S_4 = \{\not{B}, \not{B}, 2\}$ $\quad S_3 = \{\not{B}, 4, \not{B}\}$
  $S_4' = \{2\}$ $\quad S_3' = \{4\}$

Properties of Multi-Set Orderings

Property 3

Let $>$ be an ordering. Then

1. $\succ_{\text{mul}}$ is an ordering.
2. if $\succ$ well-founded then $\succ_{\text{mul}}$ well-founded.
3. if $\succ$ total then $\succ_{\text{mul}}$ total

Reconsider Example

- $S_1 = \{\not{B}, 5, \not{A}, \not{B}, \not{2}\}$ $\quad S_2 = \{\not{B}, \not{A}, 4, \not{B}, 3, \not{2}\}$
  $S_1' = \{5\}$ $\quad S_2' = \{4, 3\}$
  $5 > 4$ and $5 > 3$
  Therefore $S_1 >_{\text{mul}} S_2$.
- $S_2 = \{\not{B}, \not{A}, 4, \not{B}, 3, 2\}$ $\quad S_3 = \{\not{B}, \not{A}, \not{B}\}$
  $S_2' = \{4, 3, 2\}$ $\quad S_3' = \emptyset$
  Therefore $S_2 >_{\text{mul}} S_3$.
- Exercise: How does $S_4 = \{5, 3, 2\}$ compare with S_3?

  Answer: $S_4 = \{\not{B}, \not{B}, 2\}$ $\quad S_3 = \{\not{B}, 4, \not{B}\}$
  $S_4' = \{2\}$ $\quad S_3' = \{4\}$
  Therefore $S_3 >_{\text{mul}} S_4$.

Summary

- (strict) orderings
- well-founded orderings
- Noetherian (well-founded) induction
- multi-sets
- multi-set ordering $\succ_{\text{mul}}$

  $=$ multi-set extension of ordering $\succ$ on the elements

COMP60121: Automated Reasoning II

Lecture 2

Previously ...

- (strict) ordering
- well-founded orderings
- multi-set orderings
- multi-sets

Recall from Part I: Literals, Clauses

- Literals

$$L \quad ::= \quad A \qquad \text{(atom, positive literal)}$$
$$\quad\quad | \quad \neg A \qquad \text{(negative literal)}$$

- Clauses

$$C, D \quad ::= \quad \bot \qquad \text{(empty clause)}$$
$$\quad\quad | \quad L_1 \vee \ldots \vee L_k, \;\; k \geq 1 \qquad \text{(non-empty clause)}$$

- **Note:**
  - We assume $\vee$ is associative and commutative, repetitions matter.
  - I.e. from now on we regard clauses as multi-sets of literals and interpret, and denote, them as disjunctions.
  - Thus, $C = P \vee P \vee \neg Q$ is identical to $C' = P \vee \neg Q \vee P$. But neither C nor C' are the same as $D = P \vee \neg Q$.

Notation in Part II

- Notation:

A, B	atoms
L	literals
C, D	clauses
$\bot$	the empty clause / falsum
N	sets of clauses
F, G	formulae

- $N \models C$ means 'any model which satisfies N also satisfies C'
  What does $N \models \bot$ mean?
- $N \vdash_{Res} C$ means 'C is derivable from N using the rules of Res'
  i.e. there is a proof of C from N in the calculus Res

Notation in Part II

- Notation:

A, B	atoms
L	literals
C, D	clauses
$\bot$	the empty clause / falsum
N	sets of clauses
F, G	formulae

- $N \models C$ means 'any model which satisfies N also satisfies C'
  What does $N \models \bot$ mean? Answer: N is unsatisfiable
- $N \vdash_{Res} C$ means 'C is derivable from N using the rules of Res'
  i.e. there is a proof of C from N in the calculus Res

Unrestricted Resolution Calculus (Variation from Part I)

- **Propositional/ground resolution calculus** Res

$$\frac{C \vee A \qquad \neg A \vee D}{C \vee D} \quad \text{(resolution)}$$

$$\frac{C \vee A \vee A}{C \vee A} \quad \text{((positive !) factoring)}$$

- Terminology: resolvent for $C \vee D$; (positive) factor for $C \vee A$
  resolved atom and factored atom, resp., for A
- These are schematic inference rules; for each substitution of the schematic variables C, D, and A, respectively, by ground clauses and ground atoms we obtain an inference rule.
- Since we assume $\vee$ is associative and commutative, note that A and $\neg A$ can occur anywhere in their respective clauses.

Recap: Conversion into Conjunctive Normal Form

- The conjunctive normal form $\mathrm{CNF}(F)$ of a formula F can be computed by applying these rewrite rules:

$$\begin{array}{rcl}
F \leftrightarrow G & \Rightarrow_{\mathrm{CNF}} & (F \rightarrow G) \wedge (G \rightarrow F) \\
F \rightarrow G & \Rightarrow_{\mathrm{CNF}} & (\neg F \vee G) \\
\neg(F \vee G) & \Rightarrow_{\mathrm{CNF}} & (\neg F \wedge \neg G) \\
\neg(F \wedge G) & \Rightarrow_{\mathrm{CNF}} & (\neg F \vee \neg G) \\
\neg\neg F & \Rightarrow_{\mathrm{CNF}} & F \\
(F \wedge G) \vee H & \Rightarrow_{\mathrm{CNF}} & (F \vee H) \wedge (G \vee H)
\end{array}$$

$$\begin{array}{rcl@{\qquad}rcl}
F \wedge \top & \Rightarrow_{\mathrm{CNF}} & F & F \wedge \bot & \Rightarrow_{\mathrm{CNF}} & \bot \\
F \vee \top & \Rightarrow_{\mathrm{CNF}} & \top & F \vee \bot & \Rightarrow_{\mathrm{CNF}} & F \\
\neg\top & \Rightarrow_{\mathrm{CNF}} & \bot & \neg\bot & \Rightarrow_{\mathrm{CNF}} & \top
\end{array}$$

- The rules may be applied in any order.
- The rules are to be applied modulo associativity and commutativity of $\wedge$ and $\vee$.

Conversion into Conjunctive Normal Form (cont'd)

- The first five rules compute the negation normal form (NNF) of a formula.
- For every formula F:
  - $\models F \leftrightarrow \mathrm{CNF}(F)$
  - $\models F \leftrightarrow \mathrm{NNF}(F)$
- Conversion to CNF (and therefore clause form) may produce a formula whose size is exponential in the size of the original formula.

Relaxing the Requirements

- The goal

  "find a formula G in CNF such that: $\models F \leftrightarrow G$"

  is therefore impractical.
- For every closed f.o. formula F:
  - $\models \mathrm{Cls}(F) \rightarrow F$, but not conversely.
  - F is (un)satisfiable iff $\mathrm{Cls}(F)$ is (un)satisfiable.
- Since we can only preserve satisfiability-equivalence anyway, there is lots of room for optimisation.
- We therefore relax the requirement to

  "find a formula G in CNF such that:
  F is satisfiable iff G is satisfiable"

  and can get efficient transformations.

Structural transformation

- Assume the context is propositional logic.
- Structural transformation (or renaming) exploits (†), left-to-right:

Property 4

$$F[G] \text{ is satisfiable iff } F[Q] \wedge (Q \leftrightarrow G) \text{ is satisfiable.} \qquad (\dagger)$$

provided Q is a (fresh) propositional variable not occuring in $F[G]$.

- Use this as a rule which introduces a new propositional symbol Q for any subformula G of F. View Q as an abbreviation for G.
- We can use this rule recursively for all subformulas in the original formula (this introduces a linear number of new propositional symbols).
- Conversion of the resulting formula to CNF increases the size only by an additional constant factor (each formula $Q \leftrightarrow G$ gives rise to at most one application of the distributivity law).

Example

- $(P \vee R) \rightarrow P$ is satisfiable iff

$$\begin{array}{ll} (Q_0 \rightarrow P) & Q_0 \text{ substituted for } (P \vee R) \\ \wedge\ (Q_0 \leftrightarrow (P \vee R)) & \end{array}$$

is satisfiable iff

$$\begin{array}{ll} Q_1 & Q_1 \text{ substituted for } (Q_0 \rightarrow P) \\ \wedge\ (Q_1 \leftrightarrow (Q_0 \rightarrow P)) & \\ \wedge\ (Q_0 \leftrightarrow (P \vee R)) & \end{array}$$

is satisfiable

Exercise

- Compute the structural transformation of $\neg P \vee (R \wedge P)$ in which a new symbol is introduced for every non-literal subformula.

Exercise

- Compute the structural transformation of $\neg P \vee (R \wedge P)$ in which a new symbol is introduced for every non-literal subformula.

$$\neg P \vee (R \wedge P)$$
$$Q_0 \quad Q_1$$

Answer: Q_0

$$\wedge\ (Q_0 \leftrightarrow (\neg P \vee Q_1))$$
$$\wedge\ (Q_1 \leftrightarrow (R \wedge P))$$

Any answer equivalent modulo the use of introduced symbols with different names is correct.

Optimising Structural Transformation

- A further improvement is possible by taking the polarity of the subformula F into account.
- Assume that F contains neither $\longrightarrow$ nor $\leftrightarrow$.
- A subformula G of F has positive polarity in F, if it occurs in the scope of an even number of negation symbols.
  It has negative polarity in F, if it occurs in the scope of an odd number of negation symbols.

Formal definition of polarity

- The notion of (positive and negative) polarity can be inductively defined as follows.
  - F has positive polarity in F.
  - Suppose G is a subformula of F.
    - If $G = \neg G'$ then G' has positive (negative) polarity in F if G has negative (positive) polarity in F.
    - If $G = G_1 \star G_2$ where $\star \in \{\vee, \wedge\}$ then G_1 and G_2 have positive (negative) polarity if G has positive (negative) polarity.

Exercise

- What is the polarity of each subformula in each of the following?
  1. $\neg P \vee (R \wedge P)$
  2. $\neg P \vee \neg(R \wedge P)$

Exercise

- What is the polarity of each subformula in each of the following?
  1. $\neg P \vee (R \wedge P)$

     $\neg P \vee (R \wedge P)$
     $+- \; + \; + \; + \; +$

     $R \wedge P$ does not occur in the scope of $\neg$.

  2. $\neg P \vee \neg(R \wedge P)$

     $\neg P \vee \neg(R \wedge P)$
     $+- \; + \; +- \; - \; -$

     $R \wedge P$ occurs in the scope of one $\neg$.

Optimising Structural Transformation (cont'd)

Property 5

Let $F[G]$ be a formula containing neither $\longrightarrow$ nor $\longleftrightarrow$;
let Q be a propositional variable not occurring in $F[G]$.

1. If G has positive polarity in F, then
   $F[G]$ is satisfiable iff $F[Q] \wedge (Q \longrightarrow G)$ is satisfiable.
2. If G has negative polarity in F, then
   $F[G]$ is satisfiable iff $F[Q] \wedge (G \longrightarrow Q)$ is satisfiable.

Structural transformation has many uses and can be generalised to inlcude $\longrightarrow$ and $\longleftrightarrow$, as well as first-order logic.

Exercise

- What is the optimised structural transformation of $\neg P \vee (R \wedge P)$ in which a new symbol is introduced for every non-literal subformula (in the original formula)?

Exercise

- What is the optimised structural transformation of $\neg P \vee (R \wedge P)$ in which a new symbol is introduced for every non-literal subformula (in the original formula)?

$$\begin{array}{l} \neg P \vee (R \wedge P) \\ \quad Q_0 \quad Q_1 \\ \rightsquigarrow \; Q_0 \\ \quad \wedge\, (Q_0 \longrightarrow (\neg P \vee Q_1)) \\ \quad \wedge\, (Q_1 \longrightarrow (R \wedge P)) \end{array}$$

Exercise (cont'd)

- What is it for $\neg P \vee \neg(R \wedge P)$?

$$
\begin{array}{ll}
 & \neg P \vee \neg(R \wedge P) \\
 & \quad Q_0 \; Q_1 \quad Q_2 \\
\rightsquigarrow & Q_0 \\
 & \wedge\, (Q_0 \longrightarrow (\neg P \vee Q_1)) \\
 & \wedge\, (Q_1 \longrightarrow \neg Q_2) \\
 & \wedge\, (Q_2 \longleftarrow (R \wedge P))
\end{array}
$$

Note that when we write 'introduce new symbols for every non-literal subformula' we are referring to the subformulae of the original formula.

Summary

- language of resolution:
  - atoms
  - literals (positive & negative)
  - clauses (= multi-sets)
- calculus Res:
  - resolution & positive factoring
- optimised conversion to clause form:
  - structural transformation

COMP60121: Automated Reasoning II

Lecture 3

Previously ...

- propositional clause logic:
  atoms, literals (positive & negative), clauses (= multi-sets)
- calculus Res:
  resolution & positive factoring
- conversion to clause form:
  - optimisation using structural transformation